

# FETOL®

REGISTRATION: Q-0209-007



## Injectable solution Oxytocic

### FORMULA

Each mL contains:

Synthetic oxytocin	10 UI
Vehicle q.s.	1 mL

### Use in species



Cattle



Horses



Sheep



Goats



Dogs



Swine



Cats

### INDICATIONS FOR USE, ROUTES OF ADMINISTRATION AND TARGET SPECIES

Drug of choice for induction of labor only for medical reasons and not for convenience. Dysfunctional uterine inertia, placental retention, expulsion of uterine exudates and fetal membranes. Stimulates milk secretion.

### DOSIS:

Cattle and horses: 10 mL.  
Swine, sheep, and goats: 3–5 mL.  
Dogs and cats: 1–3 mL.

**ROUTE OF ADMINISTRATION:** Intramuscular or subcutaneous.

### WITHDRAWAL PERIOD:

Do not administer this product within 30 days before slaughter of animals intended for human consumption.

### STORAGE CONDITIONS:

Store the product in a cool place at no more than 25 °C, dry and protected from light. Keep out of reach of children.

# FETOL®

REGISTRATION: Q-0209-007



## **WARNINGS:**

Prostaglandins may potentiate the uterotonic effect of oxytocin and vice versa. Oxytocin is contraindicated in dystocic deliveries due to inadequate fetal position and/or presentation. Do not use for induction of labor when the cervix is not dilated.

Do not administer in the last third of gestation.

Do not administer to animals with cardiovascular diseases or females predisposed to uterine rupture.

Use with caution in toxemias.

Consult a Veterinarian.

Veterinary use only.

Sale requires a medical prescription.

**For exclusive use by the Veterinarian.**