

FIBROCINE®

REGISTRATION: Q-0209-053



Injectable solution

Fibrolytic.



FORMULA

Each 100 mL contains:

Thiosinamine	7 g
Sodium iodide	7 g
Sodium salicylate	7 g
Vehicle q.s.	100 mL

Use in species



Cattle



Horses



Sheep



Goats



Dogs



Swine

INDICATIONS FOR USE AND TARGET SPECIES

Specific medication to promote absorption of pathogenic fibrous tissue, chronic inflammations, ovarian fibrosis, cervical fibrosis, fibrous mastitis, and preputial phimosis in beef and dairy cattle, sheep, goats, and dogs.

DOSAGE:

Animal	Dosage
Cattle and horses:	10 mL.
Swine, sheep, and goats:	5 mL.
Dogs:	1-2 mL.

Repeat dose every 4 days until normalization of affected tissue, up to a maximum of 6 applications, at the veterinarian's discretion.

ROUTE OF ADMINISTRATION: Intramuscular

WITHDRAWAL PERIOD:

Do not use this product within 15 days before slaughter of animals intended for human consumption. Do not administer to horses intended for human consumption.

STORAGE CONDITIONS:

Store in a cool, dry place protected from light. Keep out of reach of children.

FIBROCINE®

REGISTRATION: Q-0209-053



WARNINGS:

Do not use in cats. Do not administer to horses intended for human consumption. Consult a Veterinarian. Veterinary use only. Sale requires a medical prescription.

For exclusive use by the Veterinarian.