

ANDOFLOX 2 %

Q-0209-090



Oral solution

Broad spectrum antibacterial for birds



Formula

Each ml contains:	
Enrofloxacin	50 mg
Vehicle c.b.p.	1 ml

Use in



Fattening birds,
combat and ornament.

INDICATIONS AND USES:

Broad spectrum antibacterial for fattening, fighting or domestic birds, indicated for prevention and treatment:

Digestive tract infections caused by *E. coli* and *Salmonella* spp.

Respiratory tract infections caused by *Haemophilus* spp., *Mycoplasma* spp., *Pasteurella* spp., *Staphylococcus* spp., *Streptococcus* spp., *Corynebacterium* spp., *Pseudomona* spp., and *E. coli*.

Genital tract infections caused by *Salmonella* spp., *E. coli*, and *Mycoplasma* spp.

Septicemic infections caused by *E. coli*, *Pasteurella* spp., *Salmonella* spp., *Staphylococcus* spp. and *Erysipelothrix rhusiopathiae*.

CLINICAL PHARMACOLOGY:

• DRUG KINETIC:

Enrofloxacin shows a notable bioavailability when administered parenterally exceeding 80 and 90%. It has excellent pharmacokinetic tissular penetration properties and slow elimination, thus its clinical efficiency from one to two hours after its administration. It achieves high concentrations of the active substance in tissues and bodily fluids; maintaining this level for up to 24 hours achieving a better intracellular penetration in vivo. This results in better effectiveness in the treatment of infections caused by intracellular bacteria, strictly eliminated by the liver and partially metabolized, with ciprofloxacin being its main metabolite.

• PHARMACODYNAMICS

Enrofloxacin produces an antibacterial effect at a nucleus level on the synthesis of bacterial DNA thus blocking the protein synthesis and the genetic information, thereby producing an inability for the bacteria to multiply and consequently a bactericidal effect.

DOSAGE:

1 ml for every liter of drinking water. Every 24 hours for 3 to 5 days.

In case of mycoplasmosis, continue the treatment for 5 consecutive days.

In case of salmonellosis, continue the treatment for 10b consecutive days.

Treatment may vary at the discretion of the veterinarian.

WARNINGS:

Do not administer ANDOFLOX ® 5% to birds producing eggs for human consumption.

Store at room temperature at no more than 30°C in a dry place.

Keep out of the reach of children.

Product for exclusive use in veterinarian medicine.

Sold by prescription only.

ETIOLOGIC AGENTS:

E. coli, *Salmonella* spp., *Haemophilus* spp., *Mycoplasma* spp., *Pasteurella* spp., *Staphylococcus* spp., *Streptococcus* spp., *Corynebacterium* spp., *Pseudomona* spp., *Erysipelothrix rhusiopathiae*.

ROUTE OF ADMINISTRATION

Parenteral (Intramuscular injection, subcutaneous or slow intravenous).

WITHDRAWAL PERIOD:

Salmonella spp., *E. coli*, *Fusobacterium* spp., *Proteus* spp., *Bordetella* spp., *Pasteurella* spp., *Mycoplasma* spp., *Streptococcus* spp., *Corynebacterium* spp.

PRESENTACIONES

250 ml