

LOCIÓN ANTIVERRUGAS

Q-0209-049



Topical Solution

Topical astringent



Formula

Each 100 ml contain	
Glacial acetic acid	12 g
Salicylic acid	8 g
Phenol	18 g
Excipient	100 ml

Use in



Bovine



Equine



Ovine



Caprine



Canines



Porcine



Felines

INDICATIONS AND USES:

Astringent lotion indicated to remove superficial warts, abrasions, calluses and necrotic tissue.

CLINICAL PHARMACOLOGY

• DRUG KINETIC:

Its absorption is very scarce and limited to the external skin layer, therefore it does not represent a complete metabolism.

• PHARMACODYNAMICS

Due to its astringent, cauterizing and antiseptic activity, it causes the destruction of the superficial layer of the skin or removal of necrotic tissue as it cauterizes and disinfects the application area.

DOSAGE

Ad libitum

ROUTE OF ADMINISTRATION

Wash affected zone with soap and water, make a depression around treatment area with a curve instrument, delimit application zone placing a protective gel on periphery to avoid accidental spills, apply lotion and allow it to act. A small inflammation can occur in the treated tissue which will disappear in a few days. Apply 1 or 2 times a day until complete resolution of treated tissue at the discretion of the Veterinary Doctor.

WARNING

It is a caustic product
Store at room temperature at no more than 30°C in a dry place.
Keep out of the reach of children.
For exclusive use of veterinarians.
Sold by prescription only.

PRESENTATION

100 ml